

ForPatients

by Roche

Advanced or Metastatic Esophageal Squamous Cell Carcinoma

A Study of RO7121661 and RO7247669 Compared With Nivolumab in Participants With Advanced or Metastatic Squamous Cell Carcinoma of the Esophagus

Trial Status

Active, not recruiting

Trial Runs In

18 Countries

Trial Identifier

NCT04785820 2020-004606-60
BP42772

The source of the below information is the publicly available website [ClinicalTrials.gov](https://clinicaltrials.gov). It has been summarised and edited into simpler language.

Trial Summary:

This is a Phase II, randomized, blinded, active-controlled, global, multicenter study designed to evaluate the safety and efficacy of lomvastomig and tobemstomig, compared with nivolumab, in patients with advanced or metastatic esophageal squamous-cell carcinoma (ESCC) refractory or intolerant to fluoropyrimidine- or taxane- and platinum-based regimen. Following approval of the protocol amendment version 3, recruitment into the lomvastomig arm has been stopped. The decision to stop recruitment for lomvastomig was based on strategic considerations and not based on emerging safety and/or efficacy data. The benefit/risk assessment for lomvastomig remains unchanged. The study was planned to enroll participants randomized in a 1:1:1 ratio to receive lomvastomig, tobemstomig, or nivolumab. With version 3 of the protocol, recruitment into the lomvastomig arm has stopped, and moving forward, participants will be randomized in a 1:1 ratio to receive either tobemstomig or nivolumab.

Hoffmann-La Roche

Sponsor

Phase 2

Phase

NCT04785820 2020-004606-60 BP42772

Trial Identifiers

Eligibility Criteria:

Gender

All

Age

>=18 Years

Healthy Volunteers

No