

Choroidal neovascularization Myopia

A clinical trial to look at how safe and well faricimab works compared with ranibizumab in people with choroidal neovascularization secondary to pathologic myopia

A Study to Evaluate the Efficacy and Safety of Faricimab in Patients With Choroidal Neovascularization Secondary to Pathologic Myopia

Trial Status
Active, not recruiting

Trial Runs In
11 Countries

Trial Identifier
NCT06176352 2023-506707-25-00
CR44829

The source of the below information is the publicly available website ClinicalTrials.gov. It has been summarised and edited into simpler language.

Trial Summary:

This is a Phase III, multicenter, randomized, double-masked, active comparator-controlled study evaluating the efficacy and safety of faricimab in patients with myopic choroidal neovascularization (CNV). This non-inferiority study will compare 6.0 mg faricimab versus 0.5 mg ranibizumab administered at a pro-re-nata (PRN) dosing regimen after an initial active IVT treatment administration at randomization (Day 1).

Hoffmann-La Roche
Sponsor

Phase 3
Phase

NCT06176352 2023-506707-25-00 CR44829
Trial Identifiers

Eligibility Criteria:

Gender
All

Age
>=18 Years

Healthy Volunteers
No

1. Why is the Poyang clinical trial needed?

Myopia, commonly known as short-sightedness, is caused by a larger or longer-shaped eyeball. In pathologic myopia, the walls of the eyeball can become extremely stretched and thin. The thinning in the back of the eye causes damage in the deeper layers under the retina, a layer of cells in the back of the eye that send signals to the brain. People with pathologic myopia are at risk of an eye condition called choroidal neovascularisation

ForPatients

by Roche

secondary to pathologic myopia (mCNV). In mCNV, vision loss is caused by the abnormal growth of new, leaky blood vessels at the back of the eye.

Current treatment of mCNV is with anti-vascular endothelial growth factor (anti-VEGF) therapies, such as ranibizumab. Anti VEGF medicines are given as an injection into the eye, also known as intravitreal (IVT) injections, but frequent treatments may be required and not everyone responds to treatment. Faricimab is designed to block growth of abnormal blood vessels and the leakage of fluid from them. Faricimab is approved by health authorities to treat adults with eye diseases such as diabetic macular oedema and neovascular age-related macular degeneration. Faricimab is now being investigated as treatment for mCNV. This clinical trial aims to compare the effects, good or bad, of faricimab versus ranibizumab in people with mCNV.

2. How does the Poyang clinical trial work?

This clinical trial is recruiting people with mCNV. People can take part if they have not yet been given any treatment for mCNV. People who take part in this clinical trial (participants) will be given either faricimab or ranibizumab at the start of the trial on Day 1, and then as needed for about 11 months. The clinical trial doctor will see the participants on a monthly basis. These hospital visits will include checks to see how the participant responds to the treatment and any side effects they may have. The total time of participation in the clinical trial will be about 1 year. Participants can stop trial treatment and leave the clinical trial at any time.

3. What are the main endpoints of the Poyang clinical trial?

The main clinical trial endpoint (the main results measured in the trial to see if the drug has worked) is the change from the start of treatment (baseline) in eyesight when using corrective lenses, known as 'best-corrected visual acuity' (BCVA), measured using an eye chart over Weeks 4, 8, and 12.

The other clinical trial endpoints include:

- The number of participants whose BCVA improves, stays the same or gets worse
- The number of participants who receive only 1 treatment injection by Weeks 12, 24 and 48
- The number of IVT injections participants receive by Weeks 12, 24 and 48
- The change in thickness of the eye layers at Weeks 4, 8, 12, and over time
- The change in the amount of fluid or damage to the layers beneath the retina
- The number and seriousness of side effects
- How faricimab affects the immune system

4. Who can take part in this clinical trial?

ForPatients

by Roche

People can take part in this trial if they are at least 18 years old with fluid in the layers beneath the retina (known as 'active' mCNV) and are otherwise healthy. People may not be able to take part in this trial if their eye condition is not caused by pathologic myopia, or they have another eye condition affecting their vision. People may also not take part if they regularly take some medicines, have any major illness or surgery within 1 month of having tests to see if they can join the trial, have had cancer in the last year, have certain medical conditions (such as infection or uncontrolled blood pressure), are pregnant or breastfeeding, or are planning on becoming pregnant during the trial or within 3 months of finishing treatment.

5. What treatment will participants be given in this clinical trial?

Everyone who joins this clinical trial will be placed into one of two groups randomly (like flipping a coin) and given either:

- Faricimab, as an IVT injection once every month, as needed, for up to 11 months, or
- Ranibizumab, as an IVT injection once every month, as needed, for up to 11 months

Participants will have an equal chance of being placed in either group. This is a double-masked trial, which means that neither the participant nor the clinical trial doctor can choose or know the group the participant is in, until the trial is over. This helps to prevent bias and expectations about what will happen. However, the participant's clinical trial doctor can find out which group the participant is in, if their safety is at risk. If treatment with faricimab or ranibizumab is not needed at a visit, a simulated (false) treatment procedure called 'sham' will be given. A sham procedure is designed to feel like a real injection, but instead, an empty syringe without a needle will be pressed against the numbed (anaesthetised) eye. This makes sure the trial treatment remains 'masked' to the participant.

6. Are there any risks or benefits in taking part in this clinical trial?

The safety or effectiveness of the experimental treatment (faricimab) may not be fully known at the time of the trial. Most trials involve some risks to the participant. However, it may not be greater than the risks related to routine medical care or the natural progression of the health condition. People who would like to participate will be told about any risks and benefits of taking part in the clinical trial, as well as any additional procedures, tests, or assessments they will be asked to undergo. All of these will be described in an informed consent document (a document that provides people with the information they need to decide to volunteer for the clinical trial).

Risks associated with the clinical trial Participants may have side effects (an unwanted effect of a drug or medical treatment) from the drugs or procedures in this clinical trial. Side effects can be mild to severe, even life or sight-threatening, and vary from person to person. Participants will be closely monitored during the clinical trial; safety assessments will be performed regularly.

ForPatients

by Roche

Faricimab and ranibizumab Participants will be told about the known side effects of faricimab and ranibizumab and possible side effects based on human studies or knowledge of similar drugs. Faricimab and ranibizumab will be given as IVT injections. Participants will be told about any known side effects of IVT injections.

Potential benefits associated with the clinical trial Participants' vision may or may not improve from participation in the clinical trial. Still, the information collected may help other people with similar medical conditions in the future.